

泰国茜草科少脉粗叶木植物的订正*

朱 华

(中国科学院西双版纳热带植物园, 云南 昆明 650223)

摘要: 鲜为人知的泰国少脉粗叶木 (*Lasianthus oligoneurus*) 经研究与 *L. constrictus* 为同种, 应归并为例者的异名。此外, *Lasianthus constrictus* var. *latifolius* 一变种亦无建立的必要。

关键词: 茜草科; 少脉粗叶木; 异名; 泰国

中图分类号: Q 949 **文献标识码:** A **文章编号:** 0253 - 2700(2003)05 - 0532 - 03

Notes on Species *Lasianthus oligoneurus* (Rubiaceae) from Thailand

ZHU Hua

(Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Kunming 650223, China)

Abstract: The rarely known species *L. oligoneurus* from Thailand is recognized to be conspecific with *L. constrictus* and thus reduced to a synonym of the latter. Additionally, the name *L. constrictus* var. *latifolius* is also reduced to a synonym of *L. constrictus*.

Key words: Rubiaceae; *Lasianthus oligoneurus*; Synonym; Thailand

Lasianthus Jack is a large genus in the family Rubiaceae with more than 180 species. It occurs from tropical Asia (more than 160 species) and Australia (only one species) to tropical Africa (c. 20 species) as well as tropical America (3 species) (Zhu, 2002). The greatest species diversity is found in Borneo of tropical Asia. The species of the genus occur almost exclusively in the understory of primary forest. Therefore the species distribution patterns have significant implications to the plant biogeography of tropical Asia and also to speciation patterns in tropical rain forest.

The genus is taxonomically extremely difficult. Although a revision on the genus from Thailand has been newly made (Zhu, 2001), study of the genus for the Flora Malesiana has shown some problems with this. In the former revision, it was considered that species *L. oligoneurus* K. Schum. could be conspecific with *L. constrictus* Wight. However, the synonym could not be conclusively designated then because the observation on *L. oligoneurus* K. Schum. was based only on type photos, and Schumann's original descriptions for the species was too general to trace details. With more recent study of the type specimen of the name, it is clear that *L. oligoneurus* K. Schum. is conspecific with *L. constrictus* Wight. Therefore, the name is here formally reduced to synonymy. Additionally, the name *Lasianthus constrictus* Wight var. *latifolius* Craib is also reduced to a synonym of *Lasianthus constrictus* Wight.

* Foundation item: This research was supported by grants of the National Natural Science Foundation of China (40271048) and the Yunnan Natural Science Foundation (2002C0067M)

收稿日期: 2003 - 01 - 20, 2003 - 02 - 17 **接受发表**

作者简介: 朱华 (1960 -) 男, 理学博士, 研究员, 主要从事热带植物学研究工作。

Lasianthus constrictus Wight, Calc. Journ. Nat. Hist. 6 : 515. 1846. Type : Myanmar, Griffith s. n. (holotype K).

Lasianthus oligoneurus K. Schum., Bot. Tidsskr. 24 : 340. 1902. **syn. nov.** Type : Thailand, Chantaburi, Schmidt 390 (holotype C).

Lasianthus constrictus Wight var. *latifolius* Craib, Fl. Siam. Enum. 2 (1) : 209. 1934. **syn. nov.** Type : Thailand. Pattani : Kerr 14940 (holotype K ; isotype BK).

Distribution : Thailand, Myanmar, Malaysia and Indonesia.

Taxonomic notes : Based on Schumann's description, *L. oligoneurus* has branches puberulous when young ; leaves oblong or oblong-lanceolate with shortly acuminate apices and acute bases, 12×4.5 cm, glabrous above, minutely pilose on nerves and nervules beneath ; nerves 3 – 4 ; stipules triangular, strigillose, c. 3 mm long ; cymes sessile ; flowers sessile and drupes trilocular. It agrees with *L. constrictus* in all the details except trilocular drupes. When more materials are examined, it is revealed that *L. constrictus* Wight has drupes with 1 – 4 pyrenes, which develop from a 4-locular ovary, and drupes with 3 developed pyrenes are frequent in Malesian population. Although drupes have not been seen in the present type specimen of *L. oligoneurus* (they are clearly shed from the sheet), the type exactly match *L. constrictus*. The variety *L. constrictus* var. *latifolius*, can not be separated from *L. constrictus* Wight when more materials are examined. Therefore both *L. oligoneurus* and *L. constrictus* var. *latifolius* are here reduced to synonyms of *L. constrictus*.

Specimens examined :

Thailand : Southeastern : Chon Buri, Murata *et al.* s.n. (L) ; Trat, Murata *et al.* T-17438 (L), Geesink *et al.* 6601 (L), 6259 (L). Peninsular : Nakhon Sri Thammarat, Kerr 15435 (L), Maxwell 85 – 1112 (L), Tagawa *et al.* T-5323 (L) ; Narathiwat, Larsen *et al.* 32851 (L), 32752 (L) ; Phangnga, Beusekom *et al.* 696 (L).

Malaysia : Perak : King 'coll. 6065 (L). Sabah : Clemens *et al.* 27353 (L) ; Elmer 21358 (L) ; Beaman 7794 (L) ; Nooteboom 962 (L) ; SAN 121946 (L), SAN 70884 (L). Sarawak : S 49001 (L), S 47504 (L), S 43843 (L), S 45380 (L), S 28493 (L), S 45841 (L), S 8475 (L), S 31102 (L) ; Purseglove *et al.* 4553 (L). Selangor : FRI 1957 (L), KL 1755 (L).

Singapore : Maxwell 82 – 100 (L), 78 – 70 (L).

Indonesia : Borneo : Tawao, Elmer 21358 (L) ; Kalimantan Barat, Moge *et al.* 4166 (L), Wiriadinata 3324 (L), Hallier 2884 (L) ; Teysmann 10833 (L) ; Natuna Islands, Steenis 1114 (L) ; Kalimantan Selatan, Vogel de 886 (L), Kartawinata 774 (L), Afriastini 1359 (L) ; Kalimantan Timur, Kostermans 9872 (L), Kostermans 4546 (L) ; Liang Gagang, Hallies 2004 (L) ; Kalteng, Jarvie *et al.* 5700 (L) ; Kelumpang, Rachman *et al.* s.n. (L). Java : Koorders 4074B (L), 43989B (L) ; Beum 713 (L) ; Hochreutiner 781 (L). Sumatra : Rahmat Si Toroes 3484 (L) ; Beccari 952 (L) ; Raap 538 (L) ; Korthals s.n. (L) ; Kostermans 22002 (L), Kostermans 22046 (L) ; Wilde de *et al.* 21238 (L) ; Sumatera Selatan, Kostermans *et al.* 132 (L) ; Sumatera Utara, Lorzing 16099 (L), Lorzing 14599 (L), Rahmat Si Toroes 5437 (L) ; Riau, Soepadmo 160 (L) ; Aceh, Wilde de *et al.* 18816 (L).

Brunei : Tutong, Nangkat *et al.* BRUN 15246 (L).

Acknowledgements : I thank Prof. K. Larsen and the Herbarium of University of Aarhus (AAU) and the Botanical Museum and Herbarium of Copenhagen (C) for sending types on loan. I am grateful to Prof. P. Baas, Dr. C. E. Ridsdale, Dr. M. Roos and Dr. Ding Hou for their help and for using Herbarium, Library and research facilities of the National Herbarium of the Netherlands (NHN), and also grateful to Dr. C. M. Taylor for her help with my work on the genus. Finally I thank two anonymous reviewers for their constructive comments on this article.

Reference :

Zhu H, 2001. A taxonomic revision of the genus *Lasianthus* Jack (Rubiaceae) from Thailand [J]. *Acta Phytotax Sin* (植物分类学报), 39 (2): 116—150

Zhu H, 2002. A revision of the genus *Lasianthus* (Rubiaceae) from China [J]. *Syst Geogr Plant*, 72: 63—110

* * * * *

《云南植物研究》简介

《云南植物研究》是国家科委（79）国科发条字 341 号文批准创办的植物学专业学报，是中国科学院主管的全国性自然科学期刊，现已成为我国植物科学研究发表论文的主要学术性刊物之一，为中国生物学类科技核心期刊，中国自然科学核心期刊。本刊荣获中科院优秀期刊二等奖（1996）及一等奖（2000）、第二届全国优秀期刊三等奖（1997）及云南省优秀科技期刊一等奖（1997）等，2001 年进入国家期刊方阵，并入选国家“双效期刊”。所发表的论文被 CA、BA、CABS、中国生物学文摘、中国林业文摘、中国药学文摘、中国科学引文数据库、万方数据库及中国学术期刊（光盘版）等国内外数据库所收录。凡从事植物学各分支学科研究的科研人员及各大专院校的植物学教师及学生均可订阅本刊。

本刊报道植物学各分支学科具有创新水平的原始研究论文和简报，植物学领域的新发现及重大应用价值的新成果快报；结合本人研究工作，反映国际最新研究水平的短篇综述等。中英文稿件均受欢迎。撰稿格式请参见本刊 2003 年（25 卷 4 期）“征稿简则”要求。本刊论文发表周期 6~10 个月，热情欢迎同行学者赐稿。

《云南植物研究》为双月刊，双月 25 日出版，2004 每期定价 15 元，需要的单位或个人请到当地邮局订阅，邮发代号：64-11，若错过订阅时间的，可将款直接汇至编辑部，我部将按期定时给您邮寄。

联系地址：云南昆明市北郊黑龙潭 中国科学院昆明植物研究所
邮政编码：650204 E-mail：BianJi@mail.kib.ac.cn，YOKE@chinajournal.net.cn
Telephone：0871-5223032；FAX：0871-5223163
http://YOKE.chinajournal.net.cn http://www.kib.ac.cn/editor/